



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LUCUS A NON LUCENDO.

By A. LATHAM BAKER, Ph. D., University of Rochester, Rochester, N. Y.

I reproduce here a curious piece of mathematical legerdemain which proves without proving. It has been, with its variations, the standard for many years in most all the text books. I think it will be difficult to find a proof (?) more utterly lacking in every pedagogical requirement, and yet correct in its final conclusion.

THEOREM. *Any continuous arc of the evolute is equal to the difference between the radii of curvature of the involute that are tangent to the arc at its extremities.*

If r is the radius of curvature and a, b the coördinates of the center of curvature, then the circle of curvature is

$$(x-a)^2 + (y-b)^2 = r^2 \dots (1).$$

But for a normal through a, b we get

$$y-b = -\frac{dx}{dy}(x-a) \dots (2),$$

and since the normal is tangent to the evolute

$$y-b = \frac{db}{da}(x-a) \dots (3).$$

From (1) and (2), if we suppose the point x, y to move along the curve, and therefore y, a, b , and r to be functions of x , we get

$$(x-a)dx + (y-b)dy - (x-a)da - (y-b)db = rdr,$$

and from (2), $(x-a)dx + (y-b)dy = 0$, whence

$$(x-a)da + (y-b)db = rdr \dots (4).$$

From (3) and (4), $(x-a)\frac{da^2 + db^2}{da^2} = -rdr \dots (5).$

From (1) and (3), $(x-a)^2 \frac{da^2 + db^2}{da^2} = r^2 \dots (6).$

(5) squared, divided by (6), gives $da^2 + db^2 = dr^2 = ds^2$.

Whence $ds = \pm dr \dots (7).$

Q. E. D.

Now, to begin with, calling r the radius of curvature does not make it such. We might just as rightfully call it *not* the radius of curvature. Under this latter hypothesis what does equation (7) prove?

To call such a sequence of *disjecta membra* a proof is little short of absurdity, however correct the formal algebraic operations may be. One seldom runs across such a total lack of syllogistic sequence of thought presented under the guise of proof.

Let us see what the r of equation (7) really does represent.

(1) puts a, b at the center of a circle through x, y .

(2) puts *another* a, b on the normal through the locus of another x, y .

(3) puts a third x, y on the tangent to the locus of a, b .

(4)=1st condition + 2d condition. Unites the first and second x, y , and the first and second a, b .

(5)=4th condition + 3d condition, puts a, b on the normal to x, y , and at the center of circle r , and *tangent to the normal*, and *therefore* on the evolute, at the center of curvature, and the arbitrary r now becomes the definite R , the radius of curvature, and the a, b becomes α, β , the coördinates of the center of curvature.

(6)=1st condition + 3d condition, puts a, b at center of circle through x, y and tangent to the *secant* through x, y , and leaves r and a, b arbitrary.

(7)=5th condition + 6th condition, converts the r of (6) into the R of (5), and the a, b into α, β , and hence (7) becomes $\sqrt{(d\alpha^2 + d\beta^2)} = dS = \pm dR$, and the theorem follows.

It will be noticed that there is not a suspicion of anything pertaining to the evolute until we reach equation (5), and that then by conjunction of conditions the thing snaps into place and the a, b locus becomes, *willy nilly*, the α, β locus or evolute, and the r similarly becomes R , the radius of curvature. The terminology applied to a, b, r is ineffective. It is the conjunction of conditions that decide what they should be called.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ARITHMETIC.

142. Proposed by M. J. CRAWFORD, Principal of Crawford's Academy, Savannah, Ga.

A gentleman has a garden 400 feet long and 300 feet wide, which he wishes to raise 9 inches higher by means of the earth to be dug out of a ditch 6 feet wide and surrounding the entire garden. How deep must the ditch be?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$400 \times 300 \times \frac{9}{12} = 90000$ cubic feet of earth necessary to raise the garden.

$2(412 + 300) \times 6 = 8544$ square feet, area of surface of ditch.

$90000 \div 8544 = 10\frac{9}{78}$ feet, depth of ditch.